

# Pneumonia

Higher Percentages Are Better (some of the recommended care given to patients if appropriate\*)

| Hospital Process of Care Measure   | UNITED STATES AVERAGE | ILLINOIS AVERAGE | Percentage for ST JOHNS HOSPITAL |
|--|-----------------------|------------------|----------------------------------|
| Percent of Pneumonia Patients Given Oxygenation Assessment   | 99%                   | 99%              | 99% of 169 patients <sup>2</sup> |
| Percent of Pneumonia Patients Assessed and Given Pneumococcal Vaccination  | 81%                   | 79%              | 91% of 181 patients <sup>2</sup> |
| Percent of Pneumonia Patients Whose Initial Emergency Room Blood Culture Was Performed Prior To The Administration Of The First Hospital Dose Of Antibiotics | 90%                   | 92%              | 95% of 116 patients <sup>2</sup> |
| Percent of Pneumonia Patients Given Smoking Cessation Advice/Counseling  | 87%                   | 87%              | 90% of 84 patients <sup>2</sup>  |
| Percent of Pneumonia Patients Given Initial Antibiotic(s) within 6 Hours After Arrival   | 93%                   | 93%              | 96% of 137 patients <sup>2</sup> |
| Percent of Pneumonia Patients Given the Most Appropriate Initial Antibiotic(s)   | 87%                   | 87%              | 92% of 72 patients <sup>2</sup>  |
| Percent of Pneumonia Patients Assessed and Given Influenza Vaccination   | 79%                   | 77%              | 91% of 133 patients <sup>2</sup> |

\* The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your health care provider if you have questions about your treatment.

2: The hospital indicated that the data submitted for this measure were based on a sample of cases.

Data Last Updated: March 24, 2009